

GCE

Mathematics B MEI

H640/01: Pure Mathematics and Mechanics

A Level

Mark Scheme for June 2022

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2022

Text Instructions

1. Annotations and abbreviations

Annotation in scoris	Meaning
✓ and ✕	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
E	Explanation mark 1
SC	Special case
^	Omission sign
MR	Misread
BP	Blank page
Highlighting	
Other abbreviations in mark scheme	Meaning
E1	Mark for explaining a result or establishing a given result
dep*	Mark dependent on a previous mark, indicated by *. The * may be omitted if only previous M mark.
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working
AG	Answer given
awrt	Anything which rounds to
BC	By Calculator
DR	This indicates that the instruction In this question you must show detailed reasoning appears in the question.

2. Subject-specific Marking Instructions for AS Level Mathematics B (MEI)

- a Annotations must be used during your marking. For a response awarded zero (or full) marks a single appropriate annotation (cross, tick, M0 or ^) is sufficient, but not required.

For responses that are not awarded either 0 or full marks, you must make it clear how you have arrived at the mark you have awarded and all responses must have enough annotation for a reviewer to decide if the mark awarded is correct without having to mark it independently.

It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

Award NR (No Response)

- if there is nothing written at all in the answer space and no attempt elsewhere in the script
- OR if there is a comment which does not in any way relate to the question (e.g. 'can't do', 'don't know')
- OR if there is a mark (e.g. a dash, a question mark, a picture) which isn't an attempt at the question.

Note: Award 0 marks only for an attempt that earns no credit (including copying out the question).

If a candidate uses the answer space for one question to answer another, for example using the space for 8(b) to answer 8(a), then give benefit of doubt unless it is ambiguous for which part it is intended.

- b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct solutions leading to correct answers are awarded full marks but work must not always be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly. Correct but unfamiliar or unexpected methods are often signalled by a correct result following an apparently incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner. If you are in any doubt whatsoever you should contact your Team Leader.

- c The following types of marks are available.

M

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

A method mark may usually be implied by a correct answer unless the question includes the DR statement, the command words “Determine” or “Show that”, or some other indication that the method must be given explicitly.

A

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

B

Mark for a correct result or statement independent of Method marks.

E

A given result is to be established or a result has to be explained. This usually requires more working or explanation than the establishment of an unknown result.


Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation *isw*. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d When a part of a question has two or more ‘method’ steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation ‘dep*’ is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation FT implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only – differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, what is acceptable will be detailed in the mark scheme. If this is not the case, please escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner.
- Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be ‘follow through’. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

- f Unless units are specifically requested, there is no penalty for wrong or missing units as long as the answer is numerically correct and expressed either in SI or in the units of the question. (e.g. lengths will be assumed to be in metres unless in a particular question all the lengths are in km, when this would be assumed to be the unspecified unit.)
We are usually quite flexible about the accuracy to which the final answer is expressed; over-specification is usually only penalised where the scheme explicitly says so.
- When a value is **given** in the paper only accept an answer correct to at least as many significant figures as the given value.
 - When a value is **not given** in the paper accept any answer that agrees with the correct value to **2 s.f.** unless a different level of accuracy has been asked for in the question, or the mark scheme specifies an acceptable range.
NB for Specification A the rubric specifies 3 s.f. as standard, so this statement reads “3 s.f”
- Follow through should be used so that only one mark in any question is lost for each distinct accuracy error.
Candidates using a value of 9.80, 9.81 or 10 for g should usually be penalised for any final accuracy marks which do not agree to the value found with 9.8 which is given in the rubric.
- g Rules for replaced work and multiple attempts:
- If one attempt is clearly indicated as the one to mark, or only one is left uncrossed out, then mark that attempt and ignore the others.
 - If more than one attempt is left not crossed out, then mark the last attempt unless it only repeats part of the first attempt or is substantially less complete.
 - if a candidate crosses out all of their attempts, the assessor should attempt to mark the crossed out answer(s) as above and award marks appropriately.
- h For a genuine misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate’s data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A or B mark in the question. Marks designated as cao may be awarded as long as there are no other errors. If a candidate corrects the misread in a later part, do not continue to follow through. E marks are lost unless, by chance, the given results are established by equivalent working. Note that a miscopy of the candidate’s own working is not a misread but an accuracy error.
- i If a calculator is used, some answers may be obtained with little or no working visible. Allow full marks for correct answers provided that there is nothing in the wording of the question specifying that analytical methods are required such as the bold “In this question you must show detailed reasoning”, or the command words “Show” and “Determine. Where an answer is wrong but there is some evidence of method, allow appropriate method marks. Wrong answers with no supporting method score zero. If in doubt, consult your Team Leader.
- j If in any case the scheme operates with considerable unfairness consult your Team Leader.

Question		Answer	Marks	AO	Guidance
1	(a)	Distance $2 + 5 + 4$ $= 11$ m	M1 A1 [2]	1.1a 1.1b	adding distances in two or three sections cao
1	(b)	velocity $= \frac{5-1}{10-15}$ $v = -0.8 \text{ ms}^{-1}$	M1 A1 [2]	1.1a 1.1b	Allow sign errors but not wrong way up. Soi Oe. Mark final answer`

Question		Answer	Marks	AO	Guidance
2		$\frac{13-x}{(x-3)(x+2)} = \frac{A}{x-3} + \frac{B}{x+2}$ $13-x = A(x+2) + B(x-3)$ $A = 2, B = -3$ So $\frac{2}{x-3} - \frac{3}{x+2}$	M1 A1 A1 [3]	1.1a 1.1b 1.1b	Clearing the denominators oe For one correct coefficient Correct partial fractions seen Accept just values for A and B if defined

Question		Answer	Marks	AO	Guidance
3	(a)		<p>B1</p> <p>B1</p> <p>[2]</p>	<p>1.1b</p> <p>1.1b</p>	<p>General shape with horizontal asymptotes Allow if asymptote not drawn provided the intention is clear Must be a one-to-one function y-values $\pm \frac{\pi}{2}$ seen</p>
3	(b)	<p>DR</p> <p>Graphs intersect when $3\sin x \cos x = \cos^2 x$ Either $\cos x = 0$ giving $x = -\frac{\pi}{2}, \frac{\pi}{2}$ or $3\sin x = \cos x$ giving $\tan x = \frac{1}{3}$ $x = 0.322, x = -2.82$ to 3s.f.</p> <p>When $x = 0.322$ or $x = -2.82$ $y = 0.9$ [So the points of intersection are $(0.322, 0.9), (-2.82, 0.9), (-\frac{\pi}{2}, 0), (\frac{\pi}{2}, 0)$]</p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>[6]</p>	<p>1.1a</p> <p>1.1b</p> <p>2.1</p> <p>2.1</p> <p>2.1</p> <p>2.1</p>	<p>soi Attempt to solve $\cos x = 0$ Both values in radians needed Allow for $x = \tan^{-1} \frac{1}{3}$ SC1 award for 18.4° and -161.6° if 90° already seen Notice 0.9 is exact.</p>

		<p>Alternative method</p> <p>DR</p> <p>Graphs intersect when $3\sin x \cos x = \cos^2 x$</p> <p>Either $\cos x = 0$</p> <p>giving $x = -\frac{\pi}{2}, \frac{\pi}{2}$</p> <p>Or $3\sin x = \cos x$</p> <p>Squaring gives</p> <p>$9\sin^2 x = \cos^2 x = 1 - \sin^2 x$</p> <p>$10\sin^2 x = 1$</p> <p>$\sin x = \pm\sqrt{0.1}$</p> <p>$x = -2.820, -0.322, 0.322, 2.820$</p> <p>Select genuine roots 0.322, -2.820</p> <p>When $x = 0.322$ or $x = -2.82$ $y = 0.9$</p> <p>[So the points of intersection are $(-\frac{\pi}{2}, 0), (\frac{\pi}{2}, 0), (0.322, 0.9), (-2.820, 0.9)$]</p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>[6]</p>	<p></p> <p>1.1b</p> <p>2.1</p> <p></p> <p></p> <p>2.1</p>	<p>soi</p> <p>Attempt to solve $\cos x = 0$</p> <p>Both values in radians needed</p> <p>Complete method for finding at least one value for $\sin x$</p> <p>Both correct roots and no others in the range</p> <p>Allow awrt 0.90</p>	<p>Notice 0.9 is exact.</p>
--	--	--	---	--	---	-----------------------------

Question		Answer	Marks	AO	Guidance	
4		<p>AG</p> $\frac{1}{(2+x)^2} = \frac{1}{4\left(1+\frac{x}{2}\right)^2} = \left[\frac{1}{4}\left(1+\frac{x}{2}\right)^{-2}\right]$ $= \frac{1}{4}\left(1+(-2)\left(\frac{x}{2}\right) + \frac{(-2)(-3)}{2!}\left(\frac{x}{2}\right)^2 + \dots\right)$ $\frac{1-x}{(2+x)^2} \approx \frac{(1-x)}{4}\left(1-x+\frac{3}{4}x^2\right)$ $\approx \frac{1}{4} - \frac{1}{2}x + \frac{7}{16}x^2$	<p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p>	<p>2.1</p> <p>2.1</p> <p>2.1</p> <p>2.1</p>	<p>Dealing correctly with the 2. Need not use negative powers for this mark</p> <p>Allow for expanding $(1+kx)^{-2}$ even where the B mark is not awarded</p> <p>Attempt to multiply their expansion by the numerator</p> <p>Convincing argument</p>	<p>Note there is mathematically nothing wrong with the direct expansion</p> $(2+x)^{-2} = 2^{-2} - 2 \times 2^{-3}x + 3 \times 2^{-4}x^2$ <p>Award B1M1 if seen</p>
		<p>Alternative method</p> $\frac{1}{(2+x)^2} = \frac{3}{(2+x)^2} - \frac{1}{2+x}$ $\frac{3}{(2+x)^2} = \frac{3}{4}\left(1+(-2)\left(\frac{x}{2}\right) + \frac{(-2)(-3)}{2!}\left(\frac{x}{2}\right)^2 + \dots\right)$ $-\frac{1}{2+x} = -\frac{1}{2}\left(1-\frac{x}{2} + \frac{x^2}{4}\dots\right)$ $\approx \frac{1}{4} - \frac{1}{2}x + \frac{7}{16}x^2$	<p>M1</p> <p>B1</p> <p>M1</p> <p>A1</p>		<p>Using partial fractions – allow an arithmetic slip</p> <p>Dealing correctly with the 2. Need not use negative powers for this mark</p> <p>Allow for expanding both $(1+kx)^{-2}$ and $(1+kx)^{-1}$ even where the B mark is not awarded</p> <p>Adding terms to complete a convincing argument</p>	
			<p>[4]</p>			

Question		Answer	Marks	AO	Guidance
5	(a)	<p>Either of these is acceptable</p>	<p>B1 B1</p> <p>[2]</p>	<p>1.1b 1.1b</p>	<p>Arrows making a closed loop in roughly the right directions Tension, weight and F labelled on their triangle and 25° (or 65°) correctly labelled. May be given as a suitable angle outside the triangle.</p>
5	(b)	<p>Using the triangle of forces</p> $F = 3g \tan 25^\circ \text{ or } \text{Tension} = \frac{3g}{\cos 25^\circ}$ <p>$F = 13.7$ Tension = 32.4 N</p> <p>Alternative method Resolve vertically $T \cos 25^\circ = 3g$ $T = 32.4 \text{ N}$ Resolve horizontally $F = T \sin 25^\circ = 13.7$</p>	<p>M1 A1 A1</p> <p>M1 A1 A1</p> <p>[3]</p>	<p>1.1a 1.1b 1.1b</p>	<p>Allow sin/cos or $25^\circ/65^\circ$ interchange to find F or T cao cao Allow sin/cos interchange or $25^\circ/65^\circ$ interchange cao cao</p>

Question		Answer	Marks	AO	Guidance	
6	(a)	<ul style="list-style-type: none"> the bricks have negligible size so contact force with the plank acts at a point The mass of plank is evenly spread across its length the weight of plank acts at centre of plank. 	<p>B1</p> <p>B1</p> <p>[2]</p>	<p>3.3</p> <p>3.3</p>	<p>Allow “no size” or “size doesn’t matter” or “shape is not relevant” etc</p> <p>Allow for either statement</p> <p>Allow the plank is the same throughout or centre of mass at centre</p>	<p>Allow “weight of bricks acts on the plank at point”</p> <p>Do not allow “mass acts” at the centre</p>
6	(b)	<p>when placed at the centre tensions are equal</p> $2 \times 75 = 2.3ng + 5g$ $n = \left[\frac{101}{2.3g} \right] = 4.48... \text{ so 4 bricks}$ <p>Alternative using moments</p> $5g \times 0.4 + 2.3gn \times 0.4 = 75 \times 0.8$ $n = \left[\frac{40.4}{9.016} \right] = 4.48... \text{ so 4 bricks}$	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>[2]</p>	<p>3.1b</p> <p>1.1b</p>	<p>Using symmetry to establish equation for n soi. Allow if the weight of the plank or one of the tensions is missing</p> <p>Final answer must be the integer</p> <p>Allow if 4 seen www</p> <p>Allow for missing moment of weight or one of the tensions</p> <p>Final answer must be the integer</p> <p>Allow if 4 seen www</p>	<p>Trial and improvement may be used</p> <p>Every term must be a moment</p>
6	(c)	$2.3gnx$ Nm	<p>B1</p> <p>B1</p> <p>[2]</p>	<p>1.1b</p> <p>1.2</p>	<p>Allow positive or negative $22.54nx$</p>	<p>Allow in an equation.</p>
6	(d)	<p>4 bricks on the point of breaking x m from A</p> <p>Taking moments about A</p> $5g \times 0.4 + 4 \times 2.3gx = 75 \times 0.8$ $9.2gx = 60 - 2g$ $x = 0.448 \text{ [so 44.8 cm from A]}$	<p>M1</p> <p>A1</p> <p>A1</p>	<p>3.1b</p> <p>1.1b</p> <p>1.1b</p>	<p>Taking moments about any point to form an equation. FT their n</p> <p>All forces used in a moment. Allow sign errors. Allow an incorrect distance used. Could be an inequality</p> <p>Fully correct equation FT their n. Allow corresponding inequality</p> <p>Need not be simplified</p> <p>cao</p>	

Question	Answer	Marks	AO	Guidance
		[3]		

Question	Answer	Marks	AO	Guidance
7 (a)	$x = 14t$ $y = 7t - \frac{1}{2}gt^2 + 5$ So the position vector is $\begin{pmatrix} 14t \\ 7t - 4.9t^2 + 5 \end{pmatrix}$	B1 M1 A1 [3]	1.1b 1.1b 2.5	must be $x = \dots$ or seen as the first component of the vector. Do not award for an expression that adds a vector to a scalar allow without +5, or if -5 seen Do not award for an expression that adds a vector to a scalar Must be a single vector. Accept $\frac{1}{2}g$ in final answer Accept $14t\mathbf{i} + (7t - 4.9t^2 + 5)\mathbf{j}$ SC1 for $\begin{pmatrix} 7t - 4.9t^2 + 5 \\ 14t \end{pmatrix}$ or $\begin{pmatrix} 14t - 4.9t^2 + 5 \\ 7t \end{pmatrix}$
7 (b)	Lands when $y = 0$ $7t - 4.9t^2 + 5 = 0$ $t = 1.95$ gives $x = 14t = 27.3$ m	M1 A1 B1 [3]	3.1b 1.1b 1.1b	Award for correct quadratic or an attempt to find value of t when their quadratic $y = 0$ cao FT their t and their linear expression for x ISW where candidates find the distance from the point of projection

Question		Answer	Marks	AO	Guidance
8	(a)	$\frac{dx}{dt} = 3t^2 - 8, \quad \frac{dy}{dt} = 2t$ $\frac{dy}{dx} = \frac{2t}{3t^2 - 8}$	M1* M1 (dep) A1 [3]	1.1a 1.1b 1.1b	attempt to differentiate both parametric equations Only allow for a complete method for finding $\frac{dy}{dx}$ in terms of t using the cartesian equation of the curve Combine their derivatives to find $\frac{dy}{dx}$ Do not allow for reciprocal cao
8	(b)	AG $t^3 - 8t = 8$ and $t^2 = 4$ gives $t = -2$ $\frac{dy}{dx} = \frac{2(-2)}{3(-2)^2 - 8} = -1$ Alternative When $\frac{dy}{dx} = \frac{2t}{3t^2 - 8} = -1$ giving $3t^2 + 2t - 8 = 0$ $t = \frac{4}{3}$ or $t = -2$ When $t = -2$ the coordinates are $((-2)^3 - 8(-2), (-2)^2) = (8, 4)$ [which is P]	M1 E1 M1 E1 [2]	3.1a 1.1b	Attempt to establish the value of t at $(8, 4)$. Allow for ± 2 or 2 stated Allow for $y = 4$ used in $\frac{dy}{dx} = \frac{2\sqrt{y}}{3y - 8}$ for the M mark only AG the negativity must be clearly established from correct working Uses the value of the derivative to find the value of t at P. Allow without reference to $t = \frac{4}{3}$
8	(c)	$\frac{dy}{dx} = \frac{2t}{3t^2 - 8} = -1$ giving $3t^2 + 2t - 8 = 0$ $t = \frac{4}{3}$ or $[t = -2$ is the point P]	M1 A1 [2]	1.1a 3.2a	Equating their $\frac{dy}{dx}$ to -1 and rearranging to form quadratic equation allow www -2 need not be seen but if seen must be rejected

8	(d)	Substitute $t^2 = y$	M1	1.1a	Allow for $x^2 = t^2(t^2 - 8)^2$
		$x = t^3 - 8t \Rightarrow x^2 = t^6 - 16t^4 + 64t^2$	A1	1.1b	
		$\Rightarrow x^2 = y^3 - 16y^2 + 64y$	A1	2.1	
		Alternative method	M1		Substituting for t in their equation for x ; allow without \pm
		Substitute $t = \pm y^{\frac{1}{2}}$	A1		Soi Allow without \pm
		$x = \pm(y^{\frac{3}{2}} - 8y^{\frac{1}{2}})$	A1		must be in the form $x^2 = \dots$ from fully correct working
		$x^2 = y(y - 8)^2 = [y^3 - 16y^2 + 64y]$	A1		Need not be simplified. Do not award if \pm not seen at all
			[3]		


9	(a)	$\mathbf{a} = \frac{d\mathbf{v}}{dt} = 2kt\mathbf{i} + 6\mathbf{j}$ <p>When $t = 2$, $\mathbf{a} = 2 \times 2k\mathbf{i} + 6\mathbf{j}$</p> $ \mathbf{a} = \sqrt{(4k)^2 + 6^2} = 10$ <p>giving $16k^2 + 36 = 100$</p> <p>So $k = 2$</p>	<p>M1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>[4]</p>	<p>3.1a</p> <p>1.1b</p> <p>3.1a</p> <p>3.2a</p>	<p>differentiating the \mathbf{v} vector</p> <p>substituting $t = 2$ into their \mathbf{a} vector</p> <p>Equate the magnitude of their \mathbf{a} vector to 10</p> <p>must choose the positive value if two values seen</p>
9	(b)	$\mathbf{r} = \int \mathbf{v} dt = \frac{kt^3}{3}\mathbf{i} + 3t^2\mathbf{j} + \mathbf{c}$ <p>particle at the origin when $t = 0$ so $\mathbf{c} = \mathbf{0}$</p> $\text{So } \mathbf{r} = \frac{kt^3}{3}\mathbf{i} + 3t^2\mathbf{j} = \left[\frac{2t^3}{3}\mathbf{i} + 3t^2\mathbf{j} \right]$	<p>M1</p> <p>A1</p> <p>[2]</p>	<p>1.1a</p> <p>1.1b</p>	<p>integrating with their k or general k. Allow for a vector or for both components separately integrated.</p> <p>Condone missing $+\mathbf{c}$ or $+\mathbf{c}$ still in their answer FT their k if positive or general k used Must be in vector form</p>
9	(c)	<p>Northeast when the \mathbf{i} component = \mathbf{j} component</p> $\frac{2t^3}{3} = 3t^2$ <p>giving $t = 4.5$ s</p> <p>[$t = 0$ rejected as the particle is at the origin]</p>	<p>M1</p> <p>A1</p> <p>[2]</p>	<p>3.1b</p> <p>1.1b</p>	<p>FT their \mathbf{r}</p> <p>www</p>

Question	Answer	Marks	AO	Guidance
10	<p>Length BC: $l^2 = 30^2 + 15^2 - 2 \times 30 \times 15 \cos \theta$</p> <p>$l^2 = 1125 - 900 \cos \theta$</p> <p>$l = (1125 - 900 \cos \theta)^{\frac{1}{2}}$</p> <p>$\frac{dl}{d\theta} = \frac{1}{2}(1125 - 900 \cos \theta)^{-\frac{1}{2}} \times 900 \sin \theta$</p> <p>$\frac{d\theta}{dt} = 0.1$</p> <p>$\frac{dl}{dt} = \frac{dl}{d\theta} \times \frac{d\theta}{dt} = \frac{450 \sin \theta}{(1125 - 900 \cos \theta)^{\frac{1}{2}}} \times 0.1$</p> <p>When $\theta = \frac{\pi}{3}$</p> <p>$\frac{dl}{dt} = \frac{45 \sin \frac{\pi}{3}}{(1125 - 900 \cos \frac{\pi}{3})^{\frac{1}{2}}} = \left[\frac{45\sqrt{3}}{2 \times 15\sqrt{3}} = \frac{3}{2} \right]$</p> <p>$1.5 \text{ cm s}^{-1}$</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>[8]</p>	<p>3.1a</p> <p>1.1b</p> <p>3.1a</p> <p>1.1b</p> <p>1.2</p> <p>1.1a</p> <p>1.1a</p> <p>3.2a</p>	<p>Soi Allow equivalent in metres</p> <p>Attempt to use the chain rule</p> <p>Any form</p> <p>Soi eg from $\theta = 0.1t$</p> <p>Using the chain rule to find $\frac{dl}{dt}$</p> <p>Substitute $\theta = \frac{\pi}{3}$ into their $\frac{dl}{d\theta}$</p> <p>Must have correct unit for the value</p> <p>Allow written as cm per second oe</p> <p>If M0 awarded here allow SC1 for $BC = \sqrt{675} = 15\sqrt{3}$ found using $\theta = \frac{\pi}{3}$</p> <p>If working in metres $l^2 = 0.1125 - 0.0900 \cos \theta$</p> <p>0.015 m s⁻¹</p>

Question	Answer	Marks	AO	Guidance
	<p>Alternative method</p> $l^2 = 30^2 + 15^2 - 2 \times 30 \times 15 \cos \theta$ $l^2 = 1125 - 900 \cos \theta$ $2l \frac{dl}{d\theta} = 900 \sin \theta$ $\frac{d\theta}{dt} = 0.1$ $\frac{dl}{dt} = \frac{dl}{d\theta} \times \frac{d\theta}{dt} = \frac{450 \sin \theta}{l} \times 0.1$ <p>When $\theta = \frac{\pi}{3}$ $\frac{dl}{dt} = \frac{45 \sin \frac{\pi}{3}}{15\sqrt{3}} = \frac{3}{2}$</p> <p>1.5 cm s⁻¹</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>[8]</p>		<p>Attempt to use the implicit differentiation. Any form soi</p> <p>Using the chain rule to find $\frac{dl}{dt}$</p> <p>Substitute $\theta = \frac{\pi}{3}$ into their $\frac{dl}{d\theta}$</p> <p>Must have correct unit for the value Allow written as cm per second oe</p> <p>If working in metres = 0.1125 - 0.0900 cos θ</p> <p>0.015 m s⁻¹</p>

Question	Answer	Marks	AO	Guidance
11	<p>Let $u = 2x + k$ $2dx = du$</p> $\int \frac{2}{(2x+k)^2} dx = \int \frac{1}{u^2} du$ $= -\frac{1}{u} [+c]$ $\int_k^{2k} \frac{2}{(2x+k)^2} dx = \int_{3k}^{5k} \left(\frac{1}{u^2}\right) du = -\frac{1}{5k} + \frac{1}{3k}$ <p>Alternatively, by inspection</p> $\int_k^{2k} \frac{2}{(2x+k)^2} dx = \left[-(2x+k)^{-1} \right]_k^{2k}$ $-\frac{1}{5k} + \frac{1}{3k}$ $= \frac{2}{15k}$ <p>This is inversely proportional to k [with constant of proportionality $\frac{2}{15}$]</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>M1</p> <p>M1</p> <p>A2</p> <p>M1</p> <p>A1</p> <p>E1</p> <p>[6]</p>	<p>2.1</p> <p>2.1</p> <p>2.1</p> <p>2.1</p> <p>2.1</p> <p>2.1</p> <p>2.1</p> <p>2.2a</p>	<p>Substituting $u = 2x + k$ Allow for a different substitution giving an integral in u only Ignore limits</p> <p>Correct integrand in terms of u Ignore limits</p> <p>correct indefinite integral constant need not be seen</p> <p>substituting correct new limits into their integrated expression, or substituting in terms of x and using original limits</p> <p>Integrating by inspection to obtain any multiple of $(2x + k)^{-1}$</p> <p>Fully correct indefinite integral – need not be simplified. substituting limits into their integrated expression</p> <p>Allow $\left(-\frac{1}{5} + \frac{1}{3}\right)\frac{1}{k}$ seen</p> <p>FT their definite integral Must use phrase “inversely proportional” to k or indicates $\propto \frac{1}{k}$</p> <p>Allow for $\int \frac{a}{u^2} du$ for any constant seen</p> <p>Allow if $\frac{a}{k}$ required at the start of the argument</p>

Question	Answer	Marks	AO	Guidance
12	<p>Assume there is a prime number p which is one less than a square number</p> $p = n^2 - 1 \text{ for some positive integer } n \geq 2$ $p = (n-1)(n+1)$ <p>If $n = 2$ $p = 1 \times 3 = 3$ which is prime [$p = 2$ is not 1 less than a square number]</p> <p>If $n > 2$ then p has two [proper] factors so is not prime which is a contradiction. So there are no prime numbers other than 3 which are 1 less than a square number</p>	<p>M1*</p> <p>M1*</p> <p>E1</p> <p>E1</p> <p>(dep)</p> <p>[4]</p>	<p>2.1</p> <p>2.1</p> <p>2.1</p> <p>2.1</p>	<p>Setting up proof by contradiction</p> <p>factorising</p> <p>Considers the possibility that one factor might be 1</p> <p>Condone missing reference to $n = 2$ (or $p = 3$) for this step. Conclusion must be clear.</p> <p>Allow SC1 where M1M0 or M0M0 has been awarded and $3 = 2^2 - 1$ is established</p>

Question	Answer	Marks	AO	Guidance
13 (a)	<p>Newton's second law for the train $5 - 2 \times 0.8 = (0.5 + 0.4)a$</p> <p>Alternative $5 - 0.8 - T = 0.5a$ $T - 0.8 = 0.4a$</p> <p>giving $a = \frac{34}{9} = 3.78 \text{ m s}^{-2}$.</p> <p>Using $v = u + at$ with $u = 0, t = 1.5$</p> <p>$v = \frac{34}{9} \times 1.5 = \frac{17}{3} = 5.67 \text{ m s}^{-1}$ (3sf)</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>[4]</p>	<p>3.1b</p> <p>1.1b</p> <p>3.1b</p> <p>1.1b</p>	<p>N2L for whole train with correct mass and all forces present</p> <p>Also allow for 2 equations where both have correct mass and all forces present in each</p> <p>using <i>suvat</i> equation(s) with $u = 0$ and their $a \neq g$ leading to a value for v</p> <p>FT their a. Any form</p>
13 (b)		<p>B1</p> <p>B1</p> <p>B1</p> <p>[3]</p>	<p>1.1b</p> <p>1.1b</p> <p>1.1b</p>	<p>weights and normal reactions (must be distinct and not vertical) Allow if both components of weight given instead. Allow in addition to weight only if clear they are for working purposes only</p> <p>tensions in string and coupling parallel to inclined plane</p> <p>R marked for both parts of the train. No additional forces Allow if distinct if it is clear they are equal in later work</p>
13 (c)	<p>Newton's second law $P - 2R - 0.9g \sin 20^\circ = 0.9a$</p>	<p>M1</p> <p>A1</p> <p>[2]</p>	<p>1.1b</p> <p>3.3</p>	<p>Newton's law with $m = 0.9$. Allow for incorrect weight term(s) or R used instead of $2R$</p> <p>Fully correct Any form</p>

Question		Answer	Marks	AO	Guidance
13	(d)	<p>When $P = 5$ the equation gives $5 - 2R - 0.9g \sin 20^\circ = 0.9a$</p> <p>When $P = 5.5$ the equation gives $5.5 - 2R - 0.9g \sin 20^\circ = 0.9 \times 2a$</p> <p>Solve simultaneously giving $R = 0.742$ $\left[a = \frac{5}{9} \right]$</p> <p>Alternative method</p> <p>When $P = 5$ $a = \frac{5 - 2R - 0.9g \sin 20^\circ}{0.9}$</p> <p>When $P = 5.5$ $a_1 = \frac{5.5 - 2R - 0.9g \sin 20^\circ}{0.9}$</p> <p>So $\frac{5.5 - 2R - 0.9g \sin 20^\circ}{0.9} = 2 \left(\frac{5 - 2R - 0.9g \sin 20^\circ}{0.9} \right)$</p> <p>giving $R = 0.742$ $\left[a = \frac{5}{9} \right]$</p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>[3]</p>	<p>3.1b</p> <p>3.1b</p> <p>1.1b</p> <p></p> <p></p> <p></p> <p></p> <p></p> <p></p>	<p>establishes one equation linking R and a. FT their (c)</p> <p>establishes another equation linking R and a. Consistent with their first equation</p> <p>method need not be seen BC</p> <p>correct value for R (a is not required)</p> <p></p> <p>Finds expression for a when $P = 5$ or $P = 5.5$</p> <p>Soi</p> <p></p> <p>Links corresponding acceleration for the other value of P</p> <p>Do not allow factor of 2 on the wrong side</p> <p>correct value for R (a is not required)</p>

Question		Answer	Marks	AO	Guidance
14	(a)	<p>When $t = 0$, $82 = \theta_0 e^0$ so $\theta_0 = 82$</p> <p>$t = 5$, $27 = \theta_0 e^{-5k}$</p> <p>giving $k = \left[-\frac{1}{5} \ln \left(\frac{27}{82} \right) \right] = 0.222$ to 3 sf</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>[3]</p>	<p>3.3</p> <p>3.3</p> <p>1.1b</p>	<p>Forming an equation for k and attempt to solve</p> <p>Allow for exact value or evaluated to at least 2 s.f.</p>
14	(b)	<p>The model predicts that temperature tends to zero but if the quantity of water is small the water will warm up so it will not cool the object to zero.</p>	<p>E1</p> <p>[1]</p>	<p>3.5b</p>	<p>Must imply to the model tends to zero and this does not match the real situation.</p>
14	(c)	<p>$\ln \theta = \ln(\theta_0 e^{-kt}) = \ln \theta_0 + \ln(e^{-kt})$</p> <p>$\ln \theta = \ln 82 - 0.222t = [4.41 - 0.222t]$</p>	<p>M1</p> <p>A1</p> <p>[2]</p>	<p>2.1</p> <p>2.1</p>	<p>Taking logs and attempting to use laws of logs</p> <p>Do not award for values of a and b obtained directly from the data and the natural log form of the model.</p> <p>FT their values for θ_0 and k</p> <p>Accept as part of equation or a and b clearly stated</p>
14	(d)	<p>When $t = 0$, $\ln \theta = 3.4$</p> <p>giving $\theta = 29.96$ so 30.0°C to 3 sf</p> <p>$\theta = 29.96 e^{-0.08t}$</p> <p>$\frac{d\theta}{dt} = 29.96 \times -0.08 e^{-0.08t}$</p> <p>When $t = 0$ $\frac{d\theta}{dt} = -2.3968$</p> <p>[object is cooling by 2.4° per minute]</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>[4]</p>	<p>3.4</p> <p>3.4</p> <p>3.4</p> <p>3.4</p>	<p>Accept 30° www Must be evaluated</p> <p>Attempt to differentiate their exponential expression for θ</p> <p>Any form eg $e^{3.4} \times -0.08 e^{-0.08t}$ or $-0.08 e^{3.4 - 0.08t}$</p> <p>Allow for correct negative value for $\frac{d\theta}{dt}$ or a clear statement that the rate of cooling is 2.4° per minute. Accept $= -0.08 e^{3.4}$</p>

		<p>Alternative method When $t = 0, \ln \theta = 3.4$ giving $\theta = 29.96$ so 30.0°C to 3 sf Differentiate $\ln \theta = 3.4 - 0.08t$ w.r.t t $\frac{1}{\theta} \frac{d\theta}{dt} = -0.08$ $\frac{d\theta}{dt} = -0.08\theta$ When $t = 0, \theta = 29.96$ so $\frac{d\theta}{dt} = -2.3968$ object is cooling by 2.4° per minute</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>[4]</p>	<p>3.4</p> <p>3.4</p> <p>3.4</p> <p>3.4</p> <p>[4]</p>	<p>Accept 30° www</p> <p>Uses implicit differentiation w.r.t t</p> <p>Correct derivative</p> <p>Allow for correct negative value for $\frac{d\theta}{dt}$ or a clear statement that the rate of cooling is 2.4° per minute</p>	
14	(e)	<p>Solve simultaneously $\ln \theta = 3.4 - 0.08t$ $\ln \theta = \ln 82 - 0.222t$ gives $t = 7.089$ $t = 7.1$ [7 minutes and 5 seconds] $\ln \theta = 2.8328$ gives $\theta = 17^\circ \text{C}$ Alternative method $82e^{-0.222t} = 30e^{-0.08t}$ $\frac{82}{30} = e^{0.142t}$ $t = 7.08$ [7 minutes and 5 seconds] $\theta = 17^\circ \text{C}$</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>[3]</p>	<p>3.1b</p> <p>3.4</p> <p>3.4</p> <p>3.4</p> <p>[3]</p>	<p>Attempting to find the intersection of their (c) and the given line</p> <p>Accept awrt 7.0, 7.1 or 7.2</p> <p>Must be the value for θ</p> <p>Equate their expressions for temperature and attempts to solve for t</p> <p>Accept awrt 7.0, 7.1 or 7.2</p> <p>Cao</p>	<p>Could be BC</p>

Need to get in touch?

If you ever have any questions about OCR qualifications or services (including administration, logistics and teaching) please feel free to get in touch with our customer support centre.

Call us on

01223 553998

Alternatively, you can email us on

support@ocr.org.uk

For more information visit



ocr.org.uk/qualifications/resource-finder



ocr.org.uk



Twitter/ocrexams



/ocrexams



/company/ocr



/ocrexams



CAMBRIDGE
UNIVERSITY PRESS & ASSESSMENT

OCR is part of Cambridge University Press & Assessment, a department of the University of Cambridge.

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored. © OCR 2022 Oxford Cambridge and RSA Examinations is a Company Limited by Guarantee. Registered in England. Registered office The Triangle Building, Shaftesbury Road, Cambridge, CB2 8EA.

Registered company number 3484466. OCR is an exempt charity.

OCR operates academic and vocational qualifications regulated by Ofqual, Qualifications Wales and CCEA as listed in their qualifications registers including A Levels, GCSEs, Cambridge Technicals and Cambridge Nationals.

OCR provides resources to help you deliver our qualifications. These resources do not represent any particular teaching method we expect you to use. We update our resources regularly and aim to make sure content is accurate but please check the OCR website so that you have the most up-to-date version. OCR cannot be held responsible for any errors or omissions in these resources.

Though we make every effort to check our resources, there may be contradictions between published support and the specification, so it is important that you always use information in the latest specification. We indicate any specification changes within the document itself, change the version number and provide a summary of the changes. If you do notice a discrepancy between the specification and a resource, please [contact us](#).

Whether you already offer OCR qualifications, are new to OCR or are thinking about switching, you can request more information using our [Expression of Interest form](#).

Please [get in touch](#) if you want to discuss the accessibility of resources we offer to support you in delivering our qualifications.